



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

THE SECOND VARIATION OF A DEFINITE INTEGRAL WHEN ONE END-POINT IS VARIABLE*

BY

GILBERT AMES BLISS

The method applied in the following paper to the discussion of the second variation in the case in which one end-point is movable on a fixed curve, is closely analogous to that of WEIERSTRASS † in his treatment of the problem for fixed end-points. The difference arises from the fact that in the present case terms outside of the integral sign must be taken into consideration. As a result of the discussion the analogue of JACOBI's criterion will be derived, defining apparently in a new way the critical point ‡ for the fixed curve along which the end-point varies. The relation between the critical and conjugate points is discussed in § 4.

§ 1. *The expression for the variation of the integral.*

Consider a fixed curve D ,

$$x = f(u), \quad y = g(u),$$

and a fixed point $B(x_1, y_1)$. Let C be a curve,

$$x = \phi(t), \quad y = \psi(t),$$

cutting D at $A(u = u_0, t = t_0)$, passing through $B(t = t_1)$, and making the integral

$$I = \int_{t_0}^{t_1} F(x, y, x', y') dt$$

a minimum with respect to values of the integral taken along other curves joining D and B , and lying in a certain neighborhood of C . The following assumptions are made: §

* Presented to the Society under a slightly different title at the Ithaca meeting, August 19, 1901. Received for publication November 27, 1901.

† *Lectures on the Calculus of Variations*, 1879.

‡ The same as KNESER's "Brennpunkt." See his *Variationsrechnung*, p. 89.

§ Literal subscripts will be used to denote differentiation, partial when several variables are involved. The zero-subscript or $[_0]$ means that in the function designated $t = t_0, u = u_0$. Unaccented letters refer to D ; while accented letters refer to C .

1) The functions discussed are regular at the points considered ;

$$2) \quad [x_u^2 + y_u^2]_0 \neq 0; \quad x'^2 + y'^2 \neq 0, \text{ for } t_0 \leq t \leq t_1;$$

3) F satisfies the usual homogeneity condition

$$(1) \quad F(x, y, \kappa x', \kappa y') = \kappa F(x, y, x', y') \quad (\kappa > 0);$$

$$(4) \quad F(x_0, y_0, x'_0, y'_0) \neq 0.$$

When the integral is taken along a curve,

$$(2) \quad \bar{x} = \phi(t) + \xi(t), \quad \bar{y} = \psi(t) + \eta(t),$$

the first variation can be put into the well-known form: *

$$(3) \quad \delta I = [F_x \xi + F_y \eta]_{t_0}^{t_1} + \int_{t_0}^{t_1} [G_1 \xi + G_2 \eta] dt,$$

where

$$G_1 = F_x - \frac{d}{dt} F_{x'}, \quad G_2 = F_y - \frac{d}{dt} F_{y'}.$$

According to WEIERSTRASS† the second variation can be expressed in the form:

$$(4) \quad \delta^2 I = [R]_{t_0}^{t_1} + \int_{t_0}^{t_1} [F_1 w'^2 + F_2 w^2] dt,$$

where

$$R = L\xi^2 + 2M\xi\eta + N\eta^2, \quad w = y'\xi - x'\eta,$$

the functions F_1 , L , M , N , F_2 being defined by the following equations:

$$F_1 = \frac{1}{y'^2} F_{x'x'} = -\frac{1}{x'y'} F_{x'y'} = \frac{1}{x'^2} F_{y'y'},$$

$$L = F_{xx'} - y'y''F_1, \quad N = F_{yy'} - x'x''F_1,$$

$$(5) \quad M = F_{x'y} + x''y'F_1 = F_{xy'} + x'y''F_1,$$

$$F_2 = \frac{1}{y'^2} (F_{xx} - y''^2 F_1 - L') = -\frac{1}{x'y'} (F_{xy} + x''y''F_1 - M')$$

$$= \frac{1}{x'^2} (F_{yy} - x''^2 F_1 - N').$$

In the first place by considering variations of the curve which pass through the end-points A and B considered as fixed, the following two necessary conditions for a minimum are found:

* See KNESER, loc. cit., § 4. The arguments of F and its derivatives are always x, y, x', y' .

† WEIERSTRASS'S *Lectures*, 1879.

I. C must be an extremal* satisfying $G_1 = 0$ and $G_2 = 0$;

II. F_1 must be ≥ 0 along the arc AB of the curve C .†

In the second place consider variations which do not pass through A . Inasmuch as only a necessary condition is desired, ξ and η can be chosen in a special manner. Let ξ_0 , η_0 , ξ , η be defined by the equations:

$$(6) \quad \xi_0 = f(u_0 + \sigma) - f(u_0) = [x_u]_0 \sigma + [x_{uu}]_0 \frac{\sigma^2}{2} + \dots,$$

$$\eta_0 = g(u_0 + \sigma) - g(u_0) = [y_u]_0 \sigma + [y_{uu}]_0 \frac{\sigma^2}{2} + \dots,$$

$$(7) \quad \xi = \phi_1 \xi_0 + \phi_2 \eta_0,$$

$$\eta = \psi_1 \xi_0 + \psi_2 \eta_0,$$

where ϕ_1 , ϕ_2 , ψ_1 , ψ_2 are functions of t satisfying the relations:

$$\phi_1(t_0) = \psi_2(t_0) = 1, \quad \phi_2(t_0) = \psi_1(t_0) = 0,$$

$$\phi_1(t_1) = \phi_2(t_1) = \psi_1(t_1) = \psi_2(t_1) = 0.$$

A curve (2) constructed with ξ and η as in (7) will be said to *belong to the class \bar{C}* . It is evident that each particular curve \bar{C} cuts D when $t = t_0$, and passes through B when $t = t_1$.

For these special variations ΔI can be expressed as a power series in σ , say

$$(8) \quad \Delta I = S_1 \sigma + S_2 \frac{\sigma^2}{2} + \dots$$

S_1 and S_2 can be calculated from δI and $\delta^2 I$. From (3) and (6), since C passes through B ,

$$(9) \quad \delta I = -[F_x x_u + F_y y_u]_0 \sigma - [F_x x_{uu} + F_y y_{uu}]_0 \frac{\sigma^2}{2} + \dots,$$

and therefore from (4) and (9),

$$S_1 = -[F_x x_u + F_y y_u]_0,$$

$$S_2 = -[F_x x_{uu} + F_y y_{uu} + Lx_u^2 + 2Mx_u y_u + Ny_u^2]_0 + \int_{t_0}^{t_1} [F_1 \bar{w}'^2 + F_2 \bar{w}^2] dt,$$

where \bar{w} and \bar{w}' are the coefficients of σ in w and its derivative.

From (8) it follows that a third necessary condition for the existence of a minimum is

III.

$$S_1 = 0.$$

* E. g., see KNESER, loc. cit., § 8.

† WEIERSTRASS'S *Lectures*, 1879.

This is the well-known condition for transversality.* It follows also from (8) that *if a minimum exists, S_2 must be ≥ 0 for all curves of class \bar{C} .* The further discussion of S_2 is the principal object of this paper.

§ 2. *A condition which prevents S_2 from becoming negative.*

Suppose now that C satisfies the conditions I and III, and (instead of II) the condition that F_1 is > 0 along the arc AB . Transform (4) by adding with Legendre,

$$0 = -[vw^2]_{t_0}^{t_1} + \int_{t_0}^{t_1} \frac{d(vw^2)}{dt} dt.$$

The integrand becomes a homogeneous quadratic expression in w and w' . If for $t_0 \leq t \leq t_1$, a regular function v exists satisfying the discriminant relation

$$(v) \quad v^2 - F_1(F_2 + v') = 0,$$

then $\delta^2 I$ becomes

$$(10) \quad \delta^2 I = [R - vw^2]_{t_0}^{t_1} + \int_{t_0}^{t_1} F_1 \left[w' + \frac{vw}{F_1} \right]^2 dt.$$

The integral of (v) is expressible in terms of the integral of a linear equation. For when $v = -F_1 U' / U$,

$$(U) \quad v^2 - F_1(F_2 + v') = \frac{F_1}{U} (F_1 U'' + F_1' U' - F_2 U) = 0.$$

Then

$$(11) \quad S_2 = - \left[F_x x_{uu} + F_{x'} y_{uu} + L x_u^2 + 2M x_u y_u + N y_u^2 + F_1 \bar{w}^2 \frac{U'}{U} \right]_0 + \int_{t_0}^{t_1} F_1 \left[\frac{U \bar{w}' - U' \bar{w}}{U} \right]^2 dt.$$

Assume the general integral of the differential equations $G_1 = 0$ and $G_2 = 0$, which are of the second order, to be

$$x = \phi(t, a, \beta), \quad y = \psi(t, a, \beta),$$

where a and β are arbitrary constants. Suppose that these equations represent C when $a = \beta = 0$. Then two particular integrals of (U) are †

$$\vartheta_1 = \begin{vmatrix} \phi_t & \phi_a \\ \psi_t & \psi_a \end{vmatrix}, \quad \vartheta_2 = \begin{vmatrix} \phi_t & \phi_\beta \\ \psi_t & \psi_\beta \end{vmatrix},$$

* KNESER, loc. cit., § 10.

† E. g., see WEIERSTRASS's *Lectures*.

where $\phi_i = \phi_i(t, 0, 0)$, etc. Suppose ϑ_1 and ϑ_2 to be linearly independent. Then the general integral of (U) is

$$(12) \quad U = c_1 \vartheta_1 + c_2 \vartheta_2.$$

Since ϑ_1 and ϑ_2 are linearly independent they satisfy the equation *

$$(13) \quad \vartheta_1 \vartheta_2' - \vartheta_2 \vartheta_1' = \frac{c}{F_1} \quad (c \neq 0).$$

A particular integral (12) can now be selected so that in S_2 the term outside of the integral vanishes. Put

$$(14) \quad \begin{aligned} P &= \left[\frac{F_{x'u} x_{uu} + F_{y'u} y_{uu}}{x_u^2 + y_u^2} \right]_0 + L_0 \cos^2 \delta + 2M_0 \sin \delta \cos \delta + N_0 \sin^2 \delta, \\ Q &= \left[F_1 \frac{(y'x_u - x'y_u)^2}{x_u^2 + y_u^2} \right]_0 = [F_1]_0 (x_0'^2 + y_0'^2) \sin^2 (\gamma - \delta), \end{aligned}$$

where γ and δ are the angles at A which C and D make with the x -axis. Then from (11),

$$S_2 = - \left[P + Q \frac{U'_0}{U_0} \right] [x_u^2 + y_u^2]_0 + \int_{t_0}^{t_1} F_1 \left[\frac{U \bar{w}' - U' \bar{w}}{U} \right]^2 dt.$$

KNESER has shown † that if $F \neq 0$ at A , and D cuts C transversally, then D cannot be tangent to C . Therefore Q is $\neq 0$. Since, furthermore, the equation (13) holds when $t = t_0$, c_1 and c_2 can be so determined that

$$(15) \quad P + Q \frac{U'_0}{U_0} = 0.$$

Two such values are

$$c_1 = P \vartheta_2(t_0) + Q \vartheta_2'(t_0), \quad -c_2 = P \vartheta_1(t_0) + Q \vartheta_1'(t_0).$$

If $H(t, t_0)$ denotes the particular integral of (U) formed with these constants, then

$$(16) \quad H(t, t_0) = P \Theta + Q \frac{\partial \Theta}{\partial t_0},$$

where

$$\Theta(t, t_0) = \begin{vmatrix} \vartheta_1(t) & \vartheta_2(t) \\ \vartheta_1(t_0) & \vartheta_2(t_0) \end{vmatrix}.$$

The integral H is useful in forming a function v to satisfy condition (v), at least when B is near A . For from (13) and (16), when $t = t_0$,

$$H_0 = Q \frac{c}{[F_1]_0} \neq 0.$$

* See CRAIG, *Linear Differential Equations*, vol. 1, p. 54.

† loc. cit., § 30.

These results lead to the following theorem:

If $H(t, t_0) \neq 0$ for $t_0 \leq t \leq t_1$, then for curves of class \bar{C} , S_2 can be expressed in the form

$$S_2 = \int_{t_0}^{t_1} F_1 \left[\frac{U\bar{w}' - U'\bar{w}}{U} \right]^2 dt,$$

which cannot become negative.

§ 3. The necessary condition.

By following still more closely the method of Weierstrass it can now be shown that the condition $H(t, t_0) \neq 0$ ($t_0 \leq t < t_1$) is necessary for the existence of a minimum. Suppose that this condition does not hold but that H has a zero t'_0 between t_0 and t_1 . Then, as will be proved, variations of class \bar{C} can be found which make S_2 and ΔI negative.

Integrate by parts the first term in the integrand of (4). Then

$$\delta^2 I = [R + F_1 w w']_{t_0}^{t_1} - \int_{t_0}^{t_1} w [F_1 w'' + F_1' w' - F_2 w] dt.$$

Consider the equation,

$$(U_\epsilon) \quad F_1 U'' + F_1' U' - (F_2 - \epsilon) U = 0,$$

where ϵ is a constant. From the theory of linear differential equations, an integral H_ϵ of the equation (U_ϵ) exists, depending upon ϵ for its value and having the following properties:

- 1) It is regular for $t_0 \leq t \leq t_1$;
- 2) $[H_\epsilon]_0 = H_0$, $[H'_\epsilon]_0 = H'_0$;
- 3) If $\eta > 0$ is selected arbitrarily, $\delta > 0$ can be found such that $|H_\epsilon - H| < \eta$ for $t_0 \leq t \leq t_1$, if $|\epsilon| < \delta$.

H and H' can not both be zero at t'_0 . For otherwise, since the functions involved are regular and $F_1 \neq 0$, the expansion of the left member of (U) could not be identically zero. From 3) therefore, δ can be chosen so small that when $|\epsilon| < \delta$, H_ϵ also vanishes between t_0 and t_1 , say at $t_{\epsilon 0}$.

Curves can now be chosen of class \bar{C} , such that w satisfies the equation (U_ϵ) . For example, let ξ and η be defined for $t_0 \leq t \leq t_{\epsilon 0}$ by the equations

$$(17) \quad \begin{aligned} w &= y'\xi - x'\eta = (y'_0\xi_0 - x'_0\eta_0) \frac{H_\epsilon}{H_0}, \\ x'\xi + y'\eta &= (x'_0\xi_0 + y'_0\eta_0) \frac{H_\epsilon}{H_0}; \end{aligned}$$

and for $t_{\epsilon 0} \leq t \leq t_1$, let $\xi = \eta = 0$. Then

$$(18) \quad \delta^2 I = [R - F_1 w w']_{t_0}^{t_1} + \epsilon \int_{t_0}^{t_1} w^2 dt.$$

From (9) and (18) by calculation as before, and since H satisfies (15), it follows that

$$S_2 = \epsilon \int_{t_0}^{t_{\epsilon_0}} \bar{w}^2 dt.$$

The function \bar{w} can not be identically zero unless H_{ϵ} is so; and by 3) H_{ϵ} can not vanish identically if δ is taken small enough, since H does not. Hence for certain functions ξ, η as in (7), $S_2 \neq 0$ and can be made positive or negative by taking values of ϵ opposite in sign. From § 1 therefore the arc AB can not make I a minimum.

If now the point A' defined on C by t'_0 is said to be the *critical point* for the curve D , a fourth necessary condition can be stated as follows:

IV. *If the extremal C , which passes through the fixed point B and cuts the fixed curve D transversally, is to make the integral*

$$I = \int_{t_0}^{t_1} F(x, y, x', y') dt$$

a minimum, then B must not lie beyond the critical point defined by D on C ; or analytically,

$$H(t, t_0) \neq 0 \quad \text{for} \quad t_0 \leq t < t_1.$$

§ 4. Relation between the conjugate and critical points.

The point conjugate to A is defined* by the zero t''_0 of $\Theta(t, t_0)$, which is nearest to t_0 . The functions Θ and H are both integrals of (U) of the form (12), and are linearly independent since $\Theta_0 = 0$ and $H_0 \neq 0$. By a theorem concerning linear differential equations of the second order† their zeros must separate each other, and $H = 0$ has therefore one root between t_0 and t''_0 .

The expression for H involves the curvature of D at A linearly. The curvature is

$$(19) \quad \frac{1}{r} = \frac{x_u y_{uu} - x_{uu} y_u}{[x_u^2 + y_u^2]^{\frac{3}{2}}}.$$

By differentiating (1) for κ it is found that

$$x' F_{x'} + y' F_{y'} = F.$$

* WEIERSTRASS'S *Lectures*, and KNESER, loc. cit., §§ 24, 31.

† M. BÔCHER, *An elementary proof of a theorem of Sturm*, Transactions of the American Mathematical Society, vol. 2 (1901), p. 150.

From this equation and III the values of F_x and F_y at A can be determined, and by substitution in (16) H becomes

$$(20) \quad H(t, t_0) = \left(\frac{P_1}{r} + P_2 \right) \Theta + Q \frac{\partial \Theta}{\partial t_0},$$

where

$$(21) \quad P_1 = \frac{F_0}{\sqrt{x_0'^2 + y_0'^2} \sin(\gamma - \delta)},$$

$$P_2 = L_0 \cos^2 \delta + 2M_0 \sin \delta \cos \delta + N_0 \sin^2 \delta.$$

Suppose C and A fixed, and D changeable but always transversal to C at A . Then if the expression (20) for H is put equal to zero and solved for r , the resulting function of t will express the value which the radius of curvature of D at A must have in order that t may determine the critical point for D . By the use of (13), (14) and (21) the function and its derivative are found to be

$$r = \frac{-P_1 \Theta}{P_2 \Theta + Q \frac{\partial \Theta}{\partial t_0}},$$

$$\frac{dr}{dt} = - \frac{c^2 \sqrt{x_0'^2 + y_0'^2} \frac{F_0}{F_1} \sin(\gamma - \delta)}{\left[P_2 \Theta + Q \frac{\partial \Theta}{\partial t_0} \right]^2}.$$

The denominator of r vanishes once between t_0 and t_0'' for the same reason that H does. From 4) of § 1 the derivative dr/dt is $\neq 0$ and has the sign of

$$\frac{F_0}{F_1} \sin(\gamma - \delta).$$

The radius of curvature (19) is positive when its direction is related to that of the curve D for increasing u as the $+y$ -axis is to the $+x$ -axis; otherwise it is negative.

From these results the following theorems can be stated if it is supposed that $F_0 > 0$:

1) *The critical point for a curve D which cuts the extremal C transversally at A , always lies between A and its conjugate A'' .*

2) *The position of the critical point is determined by the curvature of D at A .**

3) *If the radius of curvature of D at A is supposed to vary continuously from 0 to ∞ on the same side of D as the arc AB , and from ∞ to 0 on the*

*See KNESER, loc. cit., p. 111.

opposite side, then the critical point moves continuously from A to A'' when there is a minimum, and from A'' to A when there is a maximum.

§ 5. *Relation between the preceding results and those of Kneser.*
Sufficient conditions.

KNESER has derived a necessary condition which corresponds to IV. He shows that it is possible to find a set of extremals,*

$$(22) \quad x = \xi(t, a), \quad y = \eta(t, a),$$

each cutting D transversally when $t = t_0$, and giving C for $a = 0$. The curve D is then represented in the vicinity of C by the equations

$$x = \xi(t_0, a), \quad y = \eta(t_0, a),$$

where a is the parameter. The condition III of transversality requires that

$$[F_x \xi_a + F_y \eta_a]_0 = 0$$

for every a near zero, since each curve of the system (22) is transversal to D . This equation can be differentiated for a and the derivatives of F expressed in terms of F_1, L, M, N , from equations (5). From (14) and (20), P and Q depend only upon the curvature and direction of D , and are independent therefore of the parameter representation. It follows that for $a = 0$,

$$\frac{\partial}{\partial a} [F_x \xi_a + F_y \eta_a]_0 = \left[P + Q \frac{\Delta'(0, 0)}{\Delta(0, 0)} \right] [\xi_a^2 + \eta_a^2]_0 = 0,$$

where

$$\Delta(t, a) = \begin{vmatrix} \xi_t & \xi_a \\ \eta_t & \eta_a \end{vmatrix}.$$

$\Delta(t, 0)$ must therefore satisfy (15). It can be proved, as for ϑ_1 and ϑ_2 , that $\Delta(t, 0)$ is also an integral of (U) . Since both H and $\Delta(t, 0)$ are integrals of (U) satisfying (15) they must be linearly dependent. That is,

$$H(t, t_0) = C \Delta(t, 0), \quad C \neq 0.$$

The condition IV can therefore be restated in KNESER's form:

IV'. *If C as in IV is to make I a minimum, then a necessary condition is*

$$\Delta(t, 0) \neq 0 \text{ for } t_0 \leq t < t_1.$$

KNESER proves this condition† by discussing the case in which B coincides

* KNESER, loc. cit., §30.

† loc. cit., §25.

with the critical point A' . He shows that unless the envelope has a singular point of a particular kind at A' , there is no minimum, and so none when B lies beyond A' . The result is a stronger condition than IV' , namely,

$$(23) \quad \Delta(t, 0) \neq 0, \quad \text{for } t_0 \leq t \leq t_1.$$

But his proof does not hold if the envelope has the exceptional form mentioned.

The method given in § 3 applies when B lies beyond A' , and then includes KNESER's exceptional case. It cannot be used when B and A' coincide. For then it is not certain that the integral H_e can be made to vanish between t_0 and t_1 , and since w must vanish for $t = t_1$, the functions ξ, η cannot be constructed as in (17).

If the conditions II and IV are amended to read:

II_a. $F_1 > 0$ for points (x, y) on AB , and for any $(x', y') \neq (0, 0)$,

IV_a. $H(t, t_0) \neq 0$ for $t_0 \leq t \leq t_1$,

then $\Delta(t, 0)$ satisfies (23). According to KNESER a field can be constructed about AB , and the four conditions I, II_a, III, IV_a are sufficient conditions for the arc AB to make the integral a minimum.

THE UNIVERSITY OF MINNESOTA,
July, 1901.
